

Clock rates, clock settings and the physics of the space-time Lorentz transformation

J.H.Field

*Département de Physique Nucléaire et Corpusculaire, Université de Genève
24, quai Ernest-Ansermet CH-1211 Genève 4.*

E-mail: john.field@cern.ch

Abstract

A careful study is made of the operational meaning of the time symbols appearing in the space-time Lorentz transformation. Four distinct symbols, with different physical meanings, are needed to describe reciprocal measurements involving stationary and uniformly-moving clocks. Physical predictions concern only the observed rate of a clock as a function of its relative speed, not its setting. How the failure to make this distinction leads to the conventional predictions of spurious ‘relativity of simultaneity’ and ‘length contraction’ effects in special relativity is explained.

PACS 03.30.+p

The purpose of this paper is to discuss the physical consequences, for measurements of space and time intervals, of the Lorentz transformation^a(LT):

$$x' = \gamma[x - vt] \quad (1)$$

$$t' = \gamma[t - \frac{vx}{c^2}] \quad (2)$$

$$y' = y \quad (3)$$

$$z' = z \quad (4)$$

where $\gamma \equiv 1/\sqrt{1 - (v/c)^2}$ and c is the speed of light in vacuum. For this it is necessary to establish the exact operational correspondence between the mathematical symbols representing space-time events (x, y, z, t) and (x', y', z', t') in two inertial frames S and S' and actual clock readings and measured spatial intervals. In the LT (1)-(4) the Cartesian spatial axes of S and S' are parallel and the frame S' is in uniform motion, relative to S along the common x, x' axis, with velocity v .

In order apply the LT to any experiment, the times t and t' must be identified with those recorded by clocks at rest in either S or S'. The simplest experiment that can be conceived will use just one clock in each frame, say, C and C'. Using these clocks two different, but reciprocal, experiments may be performed. The two clocks may be viewed from S, or they may be viewed from S'. In order to describe these experiments, four physically distinct time symbols are required: τ, t', τ' and t . The symbols τ (τ')

^aThe Lorentz Transformation (1)-(4) was first given by Larmor in 1900 [1]. The same transformation appears in 1904 in Lorentz's last pre-relativity paper [2]. The latter paper was cited in 1905 by Poincaré [3] who first introduced the appellation ‘Lorentz Transformation’. Einstein independently derived the transformation in the same year [4]. See Ref. [5] for a discussion of historical priority issues

represent the times recorded by the stationary clocks C (C') as observed in S (S'), while the symbols t (t') represent the times recorded by the moving clocks C (C') as observed from S'(S).

To analyse these experiments it will be found convenient to use the invariant interval relations [6, 7] connecting τ, t' or τ', t that may be derived from the LT (1)-(4), or its inverse:

$$c^2(\Delta t')^2 - (\Delta x')^2 = c^2(\Delta \tau)^2 - (\Delta x)^2 \quad (5)$$

$$c^2(\Delta t)^2 - (\Delta x)^2 = c^2(\Delta \tau')^2 - (\Delta x')^2 \quad (6)$$

Eqn(5) describes the experiment where C' is viewed from S, so that so that the transformed events lie on the world line of C'. Since C' is at rest in S', $\Delta x' = 0$, while its equation of motion in S is: $\Delta x = v\Delta \tau$. Thus (5) may be written as

$$c^2(\Delta t')^2 = c^2(1 - \frac{v^2}{c^2})(\Delta \tau)^2$$

or

$$\Delta \tau = \gamma \Delta t' \quad (7)$$

Similarly, (6) describes the experiment where C is viewed from S', and the transformed events lie on the world line of C. In this case $\Delta x = 0$, and $\Delta x' = -v\Delta \tau'$ so that Eqn(6) gives:

$$\Delta \tau' = \gamma \Delta t \quad (8)$$

The equations (7) and (8) describe the relativistic time dilatation (TD) effect. In may be succinctly stated as follows:

$$\frac{\text{rate of moving clock}}{\text{rate of stationary clock}} = \frac{\Delta t'}{\Delta \tau} = \frac{\Delta t}{\Delta \tau'} = \frac{1}{\gamma} \quad (9)$$

This relation holds for clocks of any construction whatever, and is, as first emphasised by Einstein, of a purely kinematical nature.

In order to discuss length interval measurements in two different inertial frames, at least four clocks are needed, say, C_A and C_B , at rest in S, and $C_{A'}$ and $C_{B'}$ at rest in S'. Suppose that C_A and C_B lie along the x -axis, separated by the distance L . The separation of $C_{A'}$ and $C_{B'}$, L' , is chosen in such a way that when the x -coordinates of C_A and $C_{A'}$ coincide, so do those of C_B and $C_{B'}$. At this instant, suppose that the readings of the clocks in the frame S are: $\tau_0(C_A)$, $\tau_0(C_B)$, $t'_0(C_{A'})$ and $t'_0(C_{B'})$, with corresponding spatial coordinates:

$$x_0(C_A) = x_0(C_{A'}) = 0, \quad x_0(C_B) = x_0(C_{B'}) = L, \quad x'_0(C_{A'}) = 0, \quad x'_0(C_{B'}) = L'$$

In an abbreviated notation the LT for events on the world lines of $C_{A'}$ and $C_{B'}$ are:

$$x'_{A'} = \gamma[x_{A'} - v\Delta \tau_A] = 0 \quad (10)$$

$$\Delta t'_{A'} = \gamma[\Delta \tau_A - \frac{vx_{A'}}{c^2}] \quad (11)$$

$$x'_{B'} - L' = \gamma[x_{B'} - L - v\Delta \tau_B] = 0 \quad (12)$$

$$\Delta t'_{B'} = \gamma[\Delta \tau_B - \frac{v(x_{B'} - L)}{c^2}] \quad (13)$$

where, for example $x_{A'} \equiv x(C_{A'})$ and $\Delta\tau_A \equiv \tau(C_A) - \tau_0(C_A)$. Note that in (12) L is the value of $x_{B'}$ when $\Delta\tau_B = 0$, and is independent of the value of v . Eliminating $x_{A'}$ between (10) and (11) and $x_{B'}$ between (12) and (13) enables (10)-(13) to be written equivalently as:

$$x_{A'} = v\Delta\tau_A = \gamma\beta\Delta t'_{A'} \quad (14)$$

$$\Delta\tau_A = \gamma\Delta t'_{A'} \quad (15)$$

$$x_{B'} - L = v\Delta\tau_B = \gamma\beta\Delta t'_{B'} \quad (16)$$

$$\Delta\tau_B = \gamma\Delta t'_{B'} \quad (17)$$

Since events with clock settings $t'_0(C_{A'})$ and $t'_0(C_{B'})$ are simultaneous, and the clocks run at the same rate, then $t'(C_{A'})$ and $t'(C_{B'})$ are also simultaneous provided that:

$$t'(C_{A'}) - t'(C_{B'}) = t'_0(C_{A'}) - t'_0(C_{B'}) \quad (18)$$

or, rearranging:

$$\Delta t'_{A'} = \Delta t'_{B'} \equiv \Delta t' \quad (19)$$

Considering now simultaneous events in S' , and in virtue of the identity: $\gamma^2 - \gamma^2\beta^2 = 1$, (14),(15) and (16),(17) are pairs of parametric equations for hyperbolae in the $\Delta\tau$ versus x plane:

$$c^2(\Delta\tau_A)^2 - (x_{A'})^2 = c^2(\Delta t')^2 = c^2(\Delta\tau_B)^2 - (x_{B'} - L)^2 \quad (20)$$

Since (15), (17) and (19) require that $\Delta\tau_A = \gamma\Delta t' = \Delta\tau_B$, it follows from (20) that, at any instant in S :

$$x_{B'}(\beta) - x_{A'}(\beta) = L \quad (21)$$

where the β dependence of $x_{A'}$ and $x_{B'}$ following from (14) and (16) at fixed $\Delta t'$ is explicitly shown. Since $x \rightarrow x'$ as $\beta \rightarrow 0$, it follows from (21) that

$$x_{B'}(0) - x_{A'}(0) = x'_{B'} - x'_{A'} \equiv L' = L \quad (22)$$

The spatial separation of the clocks is therefore a Lorentz invariant quantity that has the same value in all inertial frames –there is no ‘relativistic length contraction’ (LC). How this spurious effect arises in the standard interpretation, in special relativity, of the LT will be explained below.

In the calculations above, no particular clock synchronisation procedure was introduced. Similar results will now be rederived using the LT with a particular choice of spatial coordinates and ‘clock synchronisation convention’. The latter concept will be first explained. A clock or a watch is intended to record time intervals. Its precision depends on how well the rate of the clock stays constant as a function of elapsed time. This requires understanding and control of the physical processes underlying the operation of the clock. Clocks and watches usually have a knob by which the actual time indicated may be ‘set’. This setting process may be performed in an arbitrary manner by the clock owner, but is completely unrelated to the physics of the clock mechanism which determines only the *rate* at which the clock runs. Sometimes, as in the case of a stopwatch, there may also be a button that simply resets the indicated time to zero. As discussed above, when the LT is used to predict the results of space-time experiments, at least one moving and one stationary clock is required. In general, the initial times displayed by

these clocks can be set to arbitrary values without affecting any physical predictions. For concreteness, suppose that both clocks are stopped (i.e. no longer record time) and the moving and stationary ones are set to the values t'_0 and τ_0 respectively. As previously, the moving clock is confined to the x -axis of the frame S. When the x -coordinates of the clocks coincide they are both started (i.e. begin to record time). The two numbers t'_0 and τ_0 define a particular ‘clock synchronisation convention’(CSC). All predictions for physical effects are independent of this convention. This method to synchronise two clocks in different inertial frames was introduced by Einstein in the original special relativity paper [4] and was termed ‘external synchronisation’ by Mansouri and Sexl [8].

It is easily seen that the LT of Eqns (1) and (2) above, cannot be used to implement a general CSC as described above. Setting $x = x' = 0$, when the clocks in S and S' have the same x -coordinate, then (1) gives $t \equiv \tau = 0$, so that, in consequence, (2) gives $t' = 0$. This means that the particular CSC: $t'_0 = \tau_0 = 0$ is built into the LT (1)-(2) at $x = x' = 0$. It is like a stop watch that can be reset, but not set! Worse, as will be shown below, it can only be reset at one particular spatial location. The modification of the LT (1)-(2) to enable the implementation of a general CSC (allowing it to describe, not only a stopwatch, but also a watch with knobs that enable to set the clock hands at any chosen positions) is straightforward. Eqns(1) and (2) are replaced by:

$$x' = \gamma[x - v(\tau - \tau_0)] \quad (23)$$

$$t' - t'_0 = \gamma[\tau - \tau_0 - \frac{vx}{c^2}] \quad (24)$$

where, as discussed above, the symbol τ is used for the time indicated by the clock at rest in S, and t' for the time indicated by the clock at rest in S', as viewed for S.

The importance of introducing the time offsets t'_0 and τ_0 to correctly describe synchronised clocks at different spatial positions (equivalent to placing additive constants on the right sides of Eqns(1) and (2)) was already pointed out by Einstein in the original 1905 special relativity paper^b:

If no assumption whatever be made as to the initial position of the moving system and as to the zero point of τ , (t' in the notation used above) an additive constant is to be placed on the right side of each of these (the LT (1)-(4))equations.

To the present writer's best knowledge, this important remark was never taken into account by Einstein, or anybody else, before the work reported in [9].

At the expense of adding the two parameters τ_0 and t'_0 the general CSC is now applied to the clocks when their x -coordinates coincide at $x = x' = 0$. However, if the clocks are somewhere else (say at $x = x' = L$), or if it is wished to use a different coordinate system to specify the position of the clocks, but to use the same CSC, (23) and (24) will not work, any more than does the LT (1) and (2). The solution, as in going from (1)-(2) to (23)-(24), is to make the replacements $\tau \rightarrow \tau - \tau(L)$ and $t' \rightarrow t' - t'(L)$ in (23) and (24) where the constants $\tau(L)$ and $t'(L)$ are now to be chosen in such a way that the two clocks at $x = x' = L$ will have the same CSC as the clocks at $x = x' = 0$ described by

^bSee P40, P134 respectively, in the English translations in Ref. [4].

(23) and (24). For this $\tau(L)$ and $t'(L)$ must satisfy the equations:

$$L = \gamma[L + v\tau(L)] \quad (25)$$

$$t'(L) = \gamma[\tau(L) + \frac{vL}{c^2}] \quad (26)$$

The solution of (25) and (26) is

$$\tau(L) = -\frac{L}{\gamma v}(\gamma - 1) \quad (27)$$

$$t'(L) = \frac{L}{\gamma v}(\gamma - 1) = -\tau(L) \quad (28)$$

With the substitutions $\tau \rightarrow \tau - \tau(L)$, $t' \rightarrow t' - t'(L)$ (23) and (24) give

$$x' = \gamma[x - v(\tau - \tau_0 - \tau(L))] \quad (29)$$

$$t' - t'_0 - t'(L) = \gamma[\tau - \tau_0 - \tau(L) - \frac{vx}{c^2}] \quad (30)$$

or, in virtue of (27) and (28),

$$x' - L = \gamma[x - L - v(\tau - \tau_0)] = 0 \quad (31)$$

$$t' - t'_0 = \gamma[\tau - \tau_0 - \frac{v(x - L)}{c^2}] \quad (32)$$

This is the LT appropriate for a clock pair at $x = x' = L$ that has the same CSC as the clock pair at $x = x' = 0$ that is described by (23) and (24).

Introducing the relative coordinates:

$$\hat{x} \equiv x - x_{C'}(\tau = \tau_0) \quad (33)$$

$$\hat{x}' \equiv x' - x'_{C'} \quad (34)$$

where $x_{C'}$ and $x'_{C'}$ are the coordinates of C' in S and S' respectively, both (23),(24) and (31),(32) may be written as:

$$\hat{x}' = \gamma[\hat{x} - v(\tau - \tau_0)] = 0 \quad (35)$$

$$t' - t'_0 = \gamma[\tau - \tau_0 - \frac{v\hat{x}}{c^2}] \quad (36)$$

which is identical to (23) and (24), or to the original LT (1) and (2), when $t'_0 = \tau_0 = 0$. except that the origin of the coordinate system in S' always coincides with the position of the clock C', whatever its spatial position. The LT (35)-(36) has been called elsewhere a 'local' LT [9].

Using (36) to eliminate $\hat{x}$ from (35) yields:

$$\tau - \tau_0 = \gamma(t' - t'_0) \quad (37)$$

which is equivalent to Eqn(7) above. (35) and (36) then predict the same TD effect for any pair of clocks with the same CSC, independently of their spatial positions.

The equations of motion of the clocks $C_{A'}$ and $C_{B'}$ in the frame S in (35) are the same as in Galilean relativity. Since the correlated LC and relativity of simultaneity effects

do not exist (the calculational error from which they originate is explained below), the sole modification of the physics of flat space-time resulting from the LT in the present example is to replace the relation

$$\tau - \tau_0 = t' - t'_0 \quad (38)$$

expressing the absolute (Newtonian) character of time in Galilean relativity, by the universal, translationally invariant, TD relation (37).

The origin of the spurious ‘relativity of simultaneity’ and ‘length contraction’ effects of conventional special relativity theory will now be explained. The LT (1)-(2) corresponds to the CSC $\tau_0 = t'_0 = 0$ for a clock $C_{A'}$ situated at the x' -origin in S' . If exactly the same LT is applied to a clock $C_{B'}$ at $x' = L'$ at $t \equiv \tau = 0$ the following equations are obtained:

$$L' = x'(C_{B'}) = \gamma x(C_{B'}) \quad (39)$$

$$t'_0 \equiv t'(C_{B'}) = -\frac{\gamma v x(C_{B'})}{c^2} = -\frac{v L'}{c^2} \quad (40)$$

Thus the CSC for $C_{B'}$ is $\tau_0 = 0$, $t'_0 = -vL'/c^2$ instead of $\tau_0 = t'_0 = 0$ when the same LT is applied at $x' = 0$. As mentioned earlier, the resettable-to-zero stopwatch described by the LT (1)-(2) at $x' = 0$ no longer works when $x' \neq 0$. In standard special relativity theory, to date, this trivial difference in the synchronisation convention given by the LT (1)-(2) when $x' \neq 0$ has been interpreted as a real physical effect –‘relativity of simultaneity’ (RS). Introducing the clock $C_{A'}$ at $x' = 0$ for which $x(C_{A'}) = x'(C_{A'}) = 0$ at $\tau = 0$, Eqn(39) may be written as:

$$L' = x'(C_{B'}) - x'(C_{A'}) = \gamma[x(C_{B'}) - x(C_{A'})] \equiv \gamma L \quad (41)$$

This is the spurious LC effect –the distance between the clocks is predicted to be shorter by the fraction $1/\gamma$ in the frame S in which they are in motion. It is also clear by considering the motion of similarly accelerated objects in a single frame of reference [9] that the LC effect of Eqn(41) is not a genuine physical effect. Both it and the RS effect of Eqn(40) are the result of misinterpretation of the time symbols in Eqns(1) and (2). These represent clock settings which are, initially, fixed at $t = t' = 0$ by the condition $x = x' = 0$, but which may, in general, be freely chosen to have any values when $x = x' = 0$, as in Eqns(23) and (24). They are not times recorded by synchronised clocks *at any values of x and x'* , as assumed in the conventional interpretation of Eqns(40) and (41). The only *physical* prediction of the LT concerning moving clocks is the TD effect of Eqn(9). The spurious RS and LC effects are the result of a confusion between arbitrary clock settings and physical time intervals. Physics determines only the *rates* of clocks. Their *settings* are arbitrary and have no physical significance.

Even though it is clear from the above that the RS effect predicted by conventional special relativity theory results from an erroneous argument, its existence (or non-existence) is still experimentally testable. Some specific satellite-borne experiments have recently been proposed to search for the RS effect [9, 10]. At the time of writing, there is ample experimental verification of TD but none of RS or LC [9].

Acknowledgements

Discussions with, or correspondence from: G.Boas, B.Echenard, M.Grünwald, Y.Keilman, M.Kloster, B.Rothenstein and D.Utterback, are gratefully acknowledged. Although they may not (yet) agree with conclusions of this paper, their questions and critical remarks were very useful to focus attention on the essential issues to be addressed.

References

- [1] J. Larmor, ‘Aether and Matter’, (CUP, Cambridge, 1900) Ch XI, P174.
- [2] H.A.Lorentz, Proceedings of the Academy of Sciences Amsterdam, 6, 1904. Reprinted in ‘The Principle of Relativity’ (Dover, New York, 1952) P11.
- [3] H.Poincaré, Compte rendu hebdomadaires des seances de l’Academie des Sciences, Paris 1905. V140, P1504.
- [4] A.Einstein, Annalen der Physik **17**, 891 (1905).
English translation by W.Perrett and G.B.Jeffery in ‘The Principle of Relativity’ (Dover, New York, 1952) P37, or in ‘Einstein’s Miraculous Year’ (Princeton University Press, Princeton, New Jersey, 1998) P123.
- [5] C.Kittel, Am. J. Phys. **42** 726 (1974)
- [6] H.Poincaré, Rend. del Circ. Mat. di Palermo **21** 129-146, 166-175 (1906).
- [7] H.Minkowski, English translation of Address to the 50th Association of German Natural Scientists and Physicians, Cologne, 21st September 1908, in ‘The Principle of Relativity’ (Dover, New York, 1952) P75.
- [8] R.Mansouri and R.U.Sexl, Gen. Rel. Grav. **8**, 497, 515, 809 (1977).
- [9] J.H.Field, ‘The Local Space-Time Lorentz Transformation: a New Formulation of Special Relativity Compatible with Translational Invariance’, arXiv pre-print: <http://xxx.lanl.gov/abs/physics/0501043>.
- [10] J.H.Field, ‘Proposals for two Satellite-Borne Experiments to Test Relativity of Simultaneity in Special Relativity’, arXiv pre-print: <http://xxx.lanl.gov/abs/physics/physics/0509213>.